

A SIMPLE PRECONDITIONER FOR A DISCONTINUOUS GALERKIN METHOD FOR THE STOKES PROBLEM

BLANCA AYUSO DE DIOS, FRANCO BREZZI, L. DONATELLA MARINI, JINCHAO XU,
AND LUDMIL ZIKATANOV

ABSTRACT. In this paper we construct Discontinuous Galerkin approximations of the Stokes problem where the velocity field is $H(\text{div}, \Omega)$ -conforming. This implies that the velocity solution is divergence-free in the whole domain. This property can be exploited to design a simple and effective preconditioner for the final linear system.

1. INTRODUCTION

In this paper we present a preconditioning strategy for a family of discontinuous Galerkin $H(\text{div}; \Omega)$ -conforming discretizations of the Stokes problem. The methods considered here were introduced in [10] where the authors showed that the approximate velocity field is truly divergence-free. These same methods were also used in [14].

In general, the numerical discretization of the Stokes problem produces algebraic linear systems of equations of the saddle-point type. Solving such algebraic linear systems has been the subject of considerable attention from various communities and nowadays at present several different approaches can be used to solve them efficiently (see [11] and references cited therein). One widely used approach is to use a block diagonal preconditioner with two blocks: one containing the inverse or a preconditioner of the stiffness matrix of a vector Poisson discretization, and one containing the inverse of a lumped mass matrix for the pressure. The resulting preconditioned system is then solved by means of the MINRES (Minimal Residual). For this type of preconditioner, a sound mathematical theory can be developed that guarantees uniform convergence (in the asymptotic limit) with respect to the mesh size of the discretization. However, it is also generally observed that with such a preconditioner the convergence of the MINRES can be quite slow.

One possible reason for the slow convergence is related to the treatment of the divergence-free condition on the velocity field. As is well known, the question of how to satisfy such a condition at the discrete level is one of the most intricate in Finite Elements.

Herein, we consider the family of discontinuous Galerkin $\mathbf{H}(\text{div}; \Omega)$ -conforming methods introduced in [10]. When appropriate finite element spaces are selected, the resulting approximation to the velocity field turns out to be exactly divergence-free. This property can be fully exploited in designing and constructing efficient preconditioners. Indeed, by using it, we can reduce the solution of the Stokes problem to the solution of a “second-order” problem in the space $\mathbf{curl} H_0^1(\Omega)$. In this way, the incompressibility constraint is incorporated into the linear system to be solved, such that the linear system is no longer in saddle-point form.

We propose then a preconditioner for the solution of the corresponding problem in $\mathbf{curl} H_0^1(\Omega)$. This is done by means of the fictitious space [16, 17] (or auxiliary space [20, 18]) framework, and the proposed preconditioner is the solution of several elliptic problems: a vector Laplacian discretized with DG- $H(\text{div}; \Omega)$ -conforming methods, and another elliptic problem discretized with an H^1 -conforming method. The solution of such systems can then be *efficiently* computed with classical approaches, for instance the Geometric Multigrid (GMG) or Algebraic Multigrid (AMG) methods.

Throughout the paper, we use the standard notation for Sobolev spaces [1]. For a bounded domain $D \subset \mathbb{R}^2$, we denote by $H^m(D)$ the L^2 -Sobolev space of order $m \geq 0$ and by $\|\cdot\|_{m,D}$ and $|\cdot|_{m,D}$ the usual Sobolev norm and seminorm, respectively. For $m = 0$, we write $L^2(D)$ instead of $H^0(D)$. For a general summability index p , we also denote by $W^{m,p}(D)$ the usual L^p -Sobolev spaces of order $m \geq 0$ with norm $\|\cdot\|_{m,p,D}$ and seminorm $|\cdot|_{m,p,D}$. By convention, we use boldface type for the vector-valued analogues: $\mathbf{H}^m(D) = [H^m(D)]^2$, likewise, we use boldface italics for the symmetric-tensor-valued analogues: $\mathcal{H}^m(D) := [H^m(D)]_{\text{sym}}^{2 \times 2}$. $H^m(D)/\mathbb{R}$ denotes the quotient space consisting of equivalence classes of elements of $H^m(D)$ that differ by a constant; for $m = 0$ the quotient space is denoted by $L^2(D)/\mathbb{R}$. We indicate by $L_0^2(D)$ the space of the $L^2(D)$ functions with zero average over D (which is obviously isomorphic to $L^2(D)/\mathbb{R}$). We use $(\cdot, \cdot)_D$ to denote the inner product in the spaces $L^2(D)$, $\mathbf{L}^2(D)$, and $\mathcal{L}^2(D)$.

Let $\Omega \subset \mathbb{R}^2$ be a polygonal domain with boundary $\Gamma = \partial\Omega$, and let $\mathbf{f} \in [L^2(\Omega)]^2$. The Stokes equations for a viscous incompressible fluid are

$$(1.1) \quad \begin{cases} -\mathbf{div}(2\nu\boldsymbol{\varepsilon}(\mathbf{u})) + \nabla p = \mathbf{f} & \text{in } \Omega \\ \mathbf{div} \mathbf{u} = 0 & \text{in } \Omega \end{cases}$$

where, with the usual notation, $\mathbf{u}$ is the velocity field, p the pressure, ν the viscosity of the fluid, and $\boldsymbol{\varepsilon}(\mathbf{u}) \in [L^2(\Omega)]_{\text{sym}}^{2 \times 2}$ the symmetric (linearized) strain rate tensor defined by $\boldsymbol{\varepsilon}(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + (\nabla \mathbf{u})^T)$. Moreover, we assume no-slip boundary conditions

$$(1.2) \quad \mathbf{u} \cdot \mathbf{n} = 0 \quad \text{on } \Gamma,$$

together with the natural condition associated with (1.1) on the tangential component of the normal stresses

$$(1.3) \quad ((2\nu\boldsymbol{\varepsilon}(\mathbf{u}) - p\mathbf{I}) \cdot \mathbf{n}) \cdot \mathbf{t} = 0 \quad \text{on } \Gamma,$$

where $\mathbf{I}$ is the identity tensor. Note that as $(\mathbf{I} \cdot \mathbf{n}) \cdot \mathbf{t} \equiv 0$ then (1.3) is reduced to

$$(1.4) \quad (\boldsymbol{\varepsilon}(\mathbf{u}) \cdot \mathbf{n}) \cdot \mathbf{t} = 0 \quad \text{on } \Gamma.$$

When the space $\mathbf{H}_{0,n}^1(\Omega) = \{\mathbf{v} \in \mathbf{H}^1(\Omega) : \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \Gamma\}$ is introduced, the variational formulation of the Stokes problem reads: *Find $(\mathbf{u}, p) \in \mathbf{H}_{0,n}^1(\Omega) \times L^2(\Omega)/\mathbb{R}$ as the solution*

of:

$$(1.5) \quad \begin{cases} a(\mathbf{u}, \mathbf{v}) + b(\mathbf{v}, p) = \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \, dx & \forall \mathbf{v} \in \mathbf{H}_{0,n}^1(\Omega) \\ b(\mathbf{u}, q) = 0 & \forall q \in L^2(\Omega)/\mathbb{R} \end{cases}$$

where the bilinear forms are defined by

$$\begin{aligned} a(\mathbf{u}, \mathbf{v}) &:= 2\nu \int_{\Omega} \boldsymbol{\varepsilon}(\mathbf{u}) : \boldsymbol{\varepsilon}(\mathbf{v}) \, dx, & \forall \mathbf{u}, \mathbf{v} \in \mathbf{H}_{0,n}^1(\Omega), \\ b(\mathbf{v}, q) &:= - \int_{\Omega} q \operatorname{div} \mathbf{v} \, dx & \forall q \in L^2(\Omega)/\mathbb{R}, \forall \mathbf{v} \in \mathbf{H}_{0,n}^1(\Omega). \end{aligned}$$

The existence and uniqueness of the solution $(\mathbf{u}, p)$ of (1.5) are very well known (see for instance [15, 12, 13]).

2. ABSTRACT SETTING AND BASIC NOTATIONS

Let $\mathcal{T}_h$ be a shape-regular family of partitions of Ω into triangles T . We denote by h_T the diameter of T , and we set $h = \max_{T \in \mathcal{T}_h} h_T$. We also assume that the decomposition $\mathcal{T}_h$ is conforming in the sense that it does not contain any hanging nodes.

We denote by $\mathcal{E}_h$ the set of all edges and by $\mathcal{E}_h^o$ and $\mathcal{E}_h^{\partial}$ the collection of all interior and boundary edges, respectively.

For $s \geq 1$, we define

$$H^s(\mathcal{T}_h) = \{ \phi \in L^2(\Omega) \text{ , such that } \phi|_T \in H^s(T), \quad \forall T \in \mathcal{T}_h \},$$

and their analogues $\mathbf{H}^s(\mathcal{T}_h)$ and $\mathcal{H}^s(\mathcal{T}_h)$, respectively. For scalar, vector-valued, and tensor functions, we use $(\cdot, \cdot)_{\mathcal{T}_h}$ to denote the $L^2(\mathcal{T}_h)$ -inner product and $\langle \cdot, \cdot \rangle_{\mathcal{E}_h}$ to denote the $L^2(\mathcal{E}_h)$ -inner product elementwise.

The vector functions are represented column-wise. We recall the definitions of the following operators acting on vectors $\mathbf{v} \in \mathbf{H}^1(\Omega)$ and on scalar functions $\phi \in H^1(\Omega)$ as

$$(2.1) \quad \operatorname{div} \mathbf{v} = \sum_{i=1}^2 \frac{\partial v^i}{\partial x_i}, \quad \operatorname{curl} \mathbf{v} = \frac{\partial v^2}{\partial x_1} - \frac{\partial v^1}{\partial x_2}, \quad \mathbf{curl} \phi = \nabla^{\perp} \phi := \left[\frac{\partial \phi}{\partial x_2}, -\frac{\partial \phi}{\partial x_1} \right]^T.$$

And, we recall the definitions of the spaces to be used herein:

$$\mathbf{H}(\operatorname{div}; \Omega) := \{ \mathbf{v} \in \mathbf{L}^2(\Omega) : \operatorname{div} \mathbf{v} \in L^2(\Omega) \},$$

$$\mathbf{H}(\operatorname{curl}; \Omega) := \{ \mathbf{v} \in \mathbf{L}^2(\Omega) : \operatorname{curl} \mathbf{v} \in L^2(\Omega) \}.$$

$$\mathbf{H}_{0,n}(\operatorname{div}; \Omega) := \{ \mathbf{v} \in \mathbf{H}(\operatorname{div}; \Omega) : \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \Gamma \},$$

$$\mathbf{H}_{0,n}(\operatorname{div}^0; \Omega) := \{ \mathbf{v} \in \mathbf{H}_{0,n}(\operatorname{div}; \Omega) : \operatorname{div} \mathbf{v} = 0 \text{ in } \Omega \}.$$

The above spaces are Hilbert spaces with the norms

$$\begin{aligned}\|\mathbf{v}\|_{H(\operatorname{div}, \Omega)}^2 &:= \|\mathbf{v}\|_{0, \Omega}^2 + \|\operatorname{div} \mathbf{v}\|_{0, \Omega}^2 & \forall \mathbf{v} \in \mathbf{H}(\operatorname{div}; \Omega), \\ \|\mathbf{v}\|_{H(\operatorname{curl}, \Omega)}^2 &:= \|\mathbf{v}\|_{0, \Omega}^2 + \|\operatorname{curl} \mathbf{v}\|_{0, \Omega}^2 & \forall \mathbf{v} \in \mathbf{H}(\operatorname{curl}; \Omega).\end{aligned}$$

Remark 2.1. *It is worth noting that if we restrict our analysis to vectors $\mathbf{u}$ and $\mathbf{v}$ in $\mathbf{H}^1(\Omega) \cap \mathbf{H}_{0,n}(\operatorname{div}^0; \Omega)$ then problem (1.5) becomes: Find $\mathbf{u} \in \mathbf{H}^1(\Omega) \cap \mathbf{H}_{0,n}(\operatorname{div}^0; \Omega)$ as the solution of:*

$$(2.2) \quad a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \, dx \quad \forall \mathbf{v} \in \mathbf{H}^1(\Omega) \cap \mathbf{H}_{0,n}(\operatorname{div}^0; \Omega).$$

As is usual in the DG approach, we now define some trace operators. Let $e \in \mathcal{E}_h^o$ be an internal edge of $\mathcal{T}_h$ shared by two elements T^1 and T^2 , and let $\mathbf{n}^1$ ($\mathbf{n}^2$) denote the unit normal on e pointing outwards from T^1 (T^2). For a scalar function $\varphi \in H^1(\mathcal{T}_h)$, a vector field $\boldsymbol{\tau} \in \mathbf{H}^1(\mathcal{T}_h)$, or a tensor field $\boldsymbol{\tau} \in \mathcal{H}^1(\mathcal{T}_h)$ we define the average operator in the usual way (see for instance [3]), that is, on internal edges

$$\{\varphi\} = \frac{1}{2}(\varphi^1 + \varphi^2), \quad \{\mathbf{v}\} = \frac{1}{2}(\mathbf{v}^1 + \mathbf{v}^2), \quad \{\boldsymbol{\tau}\} = \frac{1}{2}(\boldsymbol{\tau}^1 + \boldsymbol{\tau}^2).$$

However, on a boundary edge, we take $\{\varphi\}$, $\{\mathbf{v}\}$, and $\{\boldsymbol{\tau}\}$ as the trace of φ , $\mathbf{v}$, and $\boldsymbol{\tau}$, respectively, on that edge.

For a scalar function $\varphi \in H^1(\mathcal{T}_h)$, the jump operator is defined as

$$[\![\varphi]\!] = \varphi^1 \mathbf{n}^1 + \varphi^2 \mathbf{n}^2 \quad \text{on } e \in \mathcal{E}_h^o, \text{ and } [\![\varphi]\!] = \varphi \mathbf{n} \quad \text{on } e \in \mathcal{E}_h^\partial$$

(where obviously $\mathbf{n}$ is the outward unit normal), so that the jump of a scalar function is a vector in the normal direction.

For a vector field $\mathbf{v} \in \mathbf{H}^1(\mathcal{T}_h)$, following, for example, [4], the jump is the symmetric matrix-valued function given on e by

$$[\![\mathbf{v}]\!] = \mathbf{v}^1 \odot \mathbf{n}^1 + \mathbf{v}^2 \odot \mathbf{n}^2 \quad \text{on } e \in \mathcal{E}_h^o, \text{ and } [\![\mathbf{v}]\!] = \mathbf{v} \odot \mathbf{n} \quad \text{on } e \in \mathcal{E}_h^\partial,$$

where $\mathbf{v} \odot \mathbf{n} = (\mathbf{v} \mathbf{n}^T + \mathbf{n} \mathbf{v}^T)/2$ is the symmetric part of the tensor product of $\mathbf{v}$ and $\mathbf{n}$. Hence, the jump of a vector-valued function is a symmetric tensor.

If we denote by $\mathbf{n}_T$ the outward unit normal to ∂T , it is easy to check that

$$(2.3) \quad \sum_{T \in \mathcal{T}_h} \int_{\partial T} \mathbf{v} \cdot \mathbf{n}_T q \, ds = \sum_{e \in \mathcal{E}_h} \int_e \{\mathbf{v}\} \cdot [\![q]\!] \, ds \quad \forall \mathbf{v} \in \mathbf{H}(\operatorname{div}; \Omega), \quad \forall q \in H^1(\mathcal{T}_h).$$

Also for $\boldsymbol{\tau} \in \mathcal{H}^1(\Omega)$ and for all $\mathbf{v} \in \mathbf{H}^1(\mathcal{T}_h)$, we have

$$(2.4) \quad \sum_{T \in \mathcal{T}_h} \int_{\partial T} (\boldsymbol{\tau} \cdot \mathbf{n}_T) \cdot \mathbf{v} \, ds = \sum_{e \in \mathcal{E}_h} \int_e \{\boldsymbol{\tau}\} : [\![\mathbf{v}]\!] \, ds.$$

2.1. Discrete Spaces: General framework. We present three choices for each of the finite element spaces $\mathbf{V}_h$ and $\mathcal{Q}_h$ to approximate velocity and pressure, respectively. For each choice, we also need an additional space $\mathcal{N}_h$ made of piecewise polynomial scalars to be used as a sort of *potentials*. We will explain the reason for doing this and the way in which to do this later on. Note, too, that we will use this space more heavily in the construction of our preconditioner. The different choices for the spaces $\mathbf{V}_h$, $\mathcal{Q}_h$, and $\mathcal{N}_h$ rely on different choices of the local polynomial spaces $\mathcal{R}(T)$, $\mathcal{S}(T)$, and $\mathcal{M}(T)$, respectively, made for each element T . Specifically, we have

$$(2.5) \quad \mathbf{V}_h := \{\mathbf{v} \in \mathbf{H}(\text{div}; \Omega) : \mathbf{v}|_T \in \mathcal{R}(T) \ \forall T \in \mathcal{T}_h, \ \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \Gamma\},$$

$$(2.6) \quad \mathcal{Q}_h := \{q \in L^2(\Omega)/\mathbb{R} : q|_T \in \mathcal{S}(T) \ \forall T \in \mathcal{T}_h\},$$

and

$$(2.7) \quad \mathcal{N}_h := \{\varphi \in H_0^1(\Omega) : \varphi|_T \in \mathcal{M}(T) \ \forall T \in \mathcal{T}_h\}.$$

The three spaces $\mathbf{V}_h$, $\mathcal{Q}_h$, and $\mathcal{N}_h$ will always be related by this exact sequences:

$$(2.8) \quad 0 \longrightarrow \mathcal{N}_h \xrightarrow{\text{curl}} \mathbf{V}_h \xrightarrow{\text{div}} \mathcal{Q}_h \longrightarrow 0.$$

It is also necessary for each operator in (2.8) to have a continuous right inverse whose norm is uniformly bounded in h . For instance, it is necessary that

$$(2.9) \quad \exists \beta > 0 \text{ s.t. } \forall h, \forall q \in \mathcal{Q}_h \exists \mathbf{v} \in \mathbf{V}_h \text{ with: } \text{div } \mathbf{v} = q \quad \text{and} \quad \|\mathbf{v}\|_{0,\Omega} \leq \frac{1}{\beta} \|q\|_{0,\Omega}.$$

Remark 2.2. *In all our examples, the pair $(\mathbf{V}_h, \mathcal{Q}_h)$ is among the classical (and very old) finite element spaces specially tailored for the approximation of the Poisson equation in mixed form. In particular, properties (2.8) and (2.9) always hold.*

2.2. Examples. We now present three examples of finite element spaces that can be used in the above framework. For each example, we specify the corresponding polynomial spaces used on each element and describe the corresponding sets of degrees of freedom. We restrict our analysis to the case of triangles; the case of parallelograms can also be considered when corresponding changes are made (see [9]).

Let us first fix the notation concerning the *spaces of polynomials*. For $m \geq 0$, we denote by $\mathbb{P}^m(T)$ the space of polynomials defined on T of degree of at most m ; the corresponding vector space is denoted by $\mathbf{P}^m(T) = (\mathbb{P}^m(T))^2$. A polynomial of degree $m \geq 3$ that vanishes throughout ∂T (hence it belongs to $H_0^1(T)$) is called a *bubble (or an H-bubble) of degree m over T* . The space of bubbles of degree m over T is denoted by $HB^m(T)$. We denote by $\mathbb{P}_{\text{hom}}^m(T)$ the space of *homogeneous* polynomials of degree m , and we denote by $\mathbf{x}^\perp$ the vector $(-x_2, x_1)$.

For $m \geq 2$,

$$(2.10) \quad \mathbb{P}_m^+(T) := \mathbb{P}^m(T) + HB^{m+1}(T)$$

And, for $m \geq 1$, we set

$$(2.11) \quad \mathbf{BDM}_m(T) := \mathbf{P}^m(T), \quad \mathbf{RT}_m(T) := \mathbf{P}^m(T) \oplus \mathbf{x} \mathbb{P}_{hom}^m(T)$$

and

$$(2.12) \quad \mathbf{TR}_m(T) := \mathbf{P}^m(T) \oplus \mathbf{x}^\perp \mathbb{P}_{hom}^m(T).$$

We also consider some generalized bubbles: a vector-valued polynomial of degree $m \geq 2$ that belongs to $\mathbf{H}_{0,n}(\text{div}, T)$ (hence whose normal component vanishes throughout ∂T) is called a *D-bubble of degree m over T* . The space of D-bubbles of degree m over T is denoted by $\mathbf{DB}^m(T)$.

All the spaces used herein are well known and widely used. They are usually referred to as *Brezzi-Douglas-Marini*, *Raviart-Thomas*, and *Rotated Raviart-Thomas* spaces, respectively.

The first example follows.

1. *Raviart-Thomas* For $k \geq 1$, we take in each T , $\mathcal{S}(T) = \mathbb{P}^k(T)$, and $\mathcal{R}(T) := \mathbf{RT}_k(T)$. The degrees of freedom in $\mathbf{RT}_k(T)$ are

$$(2.13) \quad \begin{aligned} \int_e \mathbf{u} \cdot \mathbf{n}_e q \, ds & \quad \forall e \in \partial T, \forall q \in \mathbb{P}^k(e), \\ \int_T \mathbf{u} \cdot \mathbf{p} \, dx & \quad \forall \mathbf{p} \in \mathbb{P}^{k-1}(T). \end{aligned}$$

As $\mathcal{Q}_h$ is made of discontinuous piecewise polynomials, here and in the following examples the degrees of freedom in $\mathcal{S}(T)$ can be taken in an almost arbitrary way. The corresponding pair of spaces $(\mathbf{V}_h, \mathcal{Q}_h)$ gives the classical Raviart-Thomas finite element approximation for second-order elliptic equations in mixed form, as introduced in [19]. It is well known and easy to check that the pair $(\mathbf{V}_h, \mathcal{Q}_h)$ satisfies

$$(2.14) \quad \text{div}(\mathbf{V}_h) = \mathcal{Q}_h$$

and that the property (2.9) is verified. We then take $\mathcal{M}(T) := \mathbb{P}^{k+1}(T)$, and note that

$$(2.15) \quad \mathbf{curl}(\mathcal{N}_h) \subseteq \mathbf{V}_h.$$

We also note that the operator $\mathbf{curl}$ has a continuous right inverse uniformly bounded from $\mathbf{V}_h \cap \mathbf{H}_{0,n}(\text{div}^0, \Omega)$ to $\mathcal{N}_h$; that is,

$$(2.16) \quad \exists C > 0 \text{ such that } \forall h, \forall \mathbf{v}_h \in \mathbf{V}_h \cap \mathbf{H}_{0,n}(\text{div}^0, \Omega) \quad \exists \varphi \in \mathcal{N}_h, \text{ such that } \mathbf{curl} \varphi = \mathbf{v}_h \quad \text{and} \quad \|\varphi\|_{1,\Omega} \leq C \|\mathbf{v}_h\|_{0,\Omega}.$$

2. *Brezzi-Douglas-Marini*: For $k \geq 1$, we take $\mathcal{S}(T) = \mathbb{P}^{k-1}(T)$, and $\mathcal{R}(T) = \mathbf{BDM}_k(T)$. The degrees of freedom for $\mathbf{BDM}_k(T)$ are (see [5]):

$$(2.17) \quad \begin{aligned} \int_e \mathbf{u} \cdot \mathbf{n}_e q \, ds & \quad \forall e \in \partial T, \forall q \in \mathbb{P}^k(e); \\ \int_T \mathbf{u} \cdot \mathbf{v} \, dx & \quad \forall \mathbf{v} \in \mathbf{TR}_{k-2}(T) \quad k \geq 2. \end{aligned}$$

The resulting finite element pair $(\mathbf{V}_h, \mathbf{Q}_h)$ is also commonly used for the approximation of second-order elliptic equations in mixed form introduced in [8]. Also in this case it has been established that the pair $(\mathbf{V}_h, \mathbf{Q}_h)$ verifies the properties of (2.14) and (2.9). We then take $\mathcal{M}(T) := \mathbb{P}^{k+1}(T)$, and note that (2.15) and (2.16) are also satisfied.

3. *Brezzi-Douglas-Fortin-Marini*: For $k \geq 1$, we take $\mathcal{S}(T) = \mathbb{P}^k(T)$ and $\mathcal{R}(T) = \mathbf{BDFM}_k(T)$, which can be written as $\mathbf{BDFM}_k = \mathbf{BDM}_k(T) + \mathbf{DB}_{k+1}(T)$. The degrees of freedom for $\mathbf{BDFM}_k(T)$, though similar to the previous ones, are given here:

$$(2.18) \quad \begin{aligned} \int_e \mathbf{u} \cdot \mathbf{n}_e q \, ds & \quad \forall e \in \partial T, \forall q \in \mathbb{P}^k(e); \\ \int_T \mathbf{u} \cdot \mathbf{v} \, dx & \quad \forall \mathbf{v} \in \mathbf{TR}_{k-1}(T). \end{aligned}$$

The resulting finite element pair $(\mathbf{V}_h, \mathbf{Q}_h)$ gives the triangular analogue of the element $\mathbf{BDFM}_k$ introduced in [7] for the approximation of second-order elliptic equations in mixed form. It is easy to check that the pair $(\mathbf{V}_h, \mathbf{Q}_h)$ verifies (2.14) and (2.9). We then take $\mathcal{M}(T) := \mathbb{P}_{k+1}^+(T)$, and note that (2.15) and (2.16) hold.

The three choices above are quite similar to each other, and the best choice among them generally depends on the problem and the way in which the discrete solution is to be used. We also use basic approximation properties: for instance, we recall that a constant C exists such that for all $T \in \mathcal{T}_h$ and for all $\mathbf{v}$, e.s. in $\mathbf{H}^s(T)$, an interpolant $\mathbf{v}^I \in \mathcal{R}(T)$ exists such that

$$(2.19) \quad \|\mathbf{v} - \mathbf{v}^I\|_{0,T} + h_T |\mathbf{v}^I|_{1,T} \leq Ch_T^s |\mathbf{v}|_{s,T}, \quad s \leq k+1.$$

3. THE DISCONTINUOUS GALERKIN $H(\text{DIV}; \Omega)$ -CONFORMING METHOD

To introduce our DG-approximation, we start by defining, for any $\mathbf{u}, \mathbf{v} \in \mathbf{H}^2(\mathcal{T}_h)$ and any $p, q \in L^2(\Omega)/\mathbb{R}$, the bilinear forms

$$(3.1) \quad \begin{aligned} A_h(\mathbf{u}, \mathbf{v}) &= 2\nu \left[(\boldsymbol{\varepsilon}(\mathbf{u}) : \boldsymbol{\varepsilon}(\mathbf{v}))_{\mathcal{T}_h} - \langle \{\boldsymbol{\varepsilon}(\mathbf{u})\} : \llbracket \mathbf{v} \rrbracket \rangle_{\mathcal{E}_h^o} - \langle \llbracket \mathbf{u} \rrbracket : \{\boldsymbol{\varepsilon}(\mathbf{v})\} \rangle_{\mathcal{E}_h^o} \right] \\ &\quad - 2\nu \left[\langle \boldsymbol{\varepsilon}(\mathbf{u})\mathbf{n}, (\mathbf{v} \cdot \mathbf{n})\mathbf{n} \rangle_{\mathcal{E}_h^\partial} + \langle (\mathbf{u} \cdot \mathbf{n})\mathbf{n}, \boldsymbol{\varepsilon}(\mathbf{v})\mathbf{n} \rangle_{\mathcal{E}_h^\partial} \right] \\ &\quad + 2\nu \left[\sum_{e \in \mathcal{E}_h^o} \alpha h_e^{-1} \int_e \llbracket \mathbf{u} \rrbracket : \llbracket \mathbf{v} \rrbracket \, ds + \sum_{e \in \mathcal{E}_h^\partial} \alpha h_e^{-1} \int_e (\mathbf{u} \cdot \mathbf{n})(\mathbf{v} \cdot \mathbf{n}) \, ds \right] \\ B_h(\mathbf{v}, q) &= -(q, \text{div } \mathbf{v})_{\mathcal{T}_h} \quad \forall \mathbf{v} \in \mathbf{H}^2(\mathcal{T}_h), \forall q \in L^2(\Omega)/\mathbb{R} \end{aligned}$$

where as usual α is the penalty parameter that we assume to be positive and large enough.

It is easy to check that the solution $(\mathbf{u}, p)$ of (1.5) verifies:

$$(3.2) \quad \begin{cases} A_h(\mathbf{u}, \mathbf{v}) + B_h(\mathbf{v}, p) = (\mathbf{f}, \mathbf{v}) & \forall \mathbf{v} \in \mathbf{H}^2(\mathcal{T}_h) \\ B_h(\mathbf{u}, q) = 0 & \forall q \in L^2(\Omega)/\mathbb{R}. \end{cases}$$

For a general DG approximation, we now replace the spaces $\mathbf{H}^2(\mathcal{T}_h)$ and $L^2(\Omega)/\mathbb{R}$ with the discrete ones $\mathcal{X}_h$ and $\mathcal{Q}_h$, respectively. Following [10], we choose for $(\mathcal{X}_h, \mathcal{Q}_h)$ one of the pairs $(\mathbf{V}_h, \mathcal{Q}_h)$ of the previous examples in order to get a global divergence-free approximation.

More generally, we can choose a pair $(\mathbf{V}_h, \mathcal{Q}_h)$ in order to find a third space $\mathcal{N}_h$ in such a way that (2.8), (2.14), (2.9), (2.15), and (2.16) are satisfied. This set of assumptions will come out several times in the sequel and, therefore, it is helpful to give it a special name.

Definition 3.1. *In the above setting, we say that the three spaces $(\mathbf{V}_h, \mathcal{Q}_h, \mathcal{N}_h)$ satisfy Assumption **H0** if (2.8), (2.14), (2.9), (2.15) and (2.16) are satisfied.*

We note that, according to the definition of $\mathbf{V}_h$, the normal component of any $\mathbf{v} \in \mathbf{V}_h$ is continuous on the internal edges and vanishes on the boundary edges. Therefore, by splitting a vector $\mathbf{v} \in \mathbf{V}_h$ into its tangential and normal components $\mathbf{v}_n$ and $\mathbf{v}_t$

$$(3.3) \quad \mathbf{v}_n := (\mathbf{v} \cdot \mathbf{n})\mathbf{n}, \quad \mathbf{v}_t := (\mathbf{v} \cdot \mathbf{t})\mathbf{t} \equiv \mathbf{v} - \mathbf{v}_n,$$

we have

$$(3.4) \quad \forall e \in \mathcal{E}_h \quad \int_e \llbracket \mathbf{v}_n \rrbracket : \boldsymbol{\tau} \, ds = 0 \quad \forall \boldsymbol{\tau} \in \boldsymbol{\mathcal{H}}^1(\mathcal{T}_h),$$

implying that

$$(3.5) \quad \forall e \in \mathcal{E}_h \quad \int_e \llbracket \mathbf{v} \rrbracket : \boldsymbol{\tau} \, ds = \int_e \llbracket \mathbf{v}_t \rrbracket : \boldsymbol{\tau} \, ds \quad \forall \boldsymbol{\tau} \in \boldsymbol{\mathcal{H}}^1(\mathcal{T}_h).$$

The resulting approximation to (1.5), therefore, becomes: *Find $(\mathbf{u}_h, p_h)$ in $\mathbf{V}_h \times \mathcal{Q}_h$ such that*

$$(3.6) \quad \begin{cases} a_h(\mathbf{u}_h, \mathbf{v}) + b(\mathbf{v}, p_h) = (\mathbf{f}, \mathbf{v}) & \forall \mathbf{v} \in \mathbf{V}_h \\ b(\mathbf{u}_h, q) = 0 & \forall q \in \mathcal{Q}_h, \end{cases}$$

where

$$(3.7) \quad \begin{aligned} a_h(\mathbf{u}, \mathbf{v}) &:= 2\nu \left[(\boldsymbol{\varepsilon}(\mathbf{u}) : \boldsymbol{\varepsilon}(\mathbf{v}))_{\mathcal{T}_h} - \langle \{\boldsymbol{\varepsilon}(\mathbf{u})\} : \llbracket \mathbf{v}_t \rrbracket \rangle_{\mathcal{E}_h^o} - \langle \llbracket \mathbf{u}_t \rrbracket : \{\boldsymbol{\varepsilon}(\mathbf{v})\} \rangle_{\mathcal{E}_h^o} \right] \\ &+ 2\nu\alpha \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \int_e \llbracket \mathbf{u}_t \rrbracket : \llbracket \mathbf{v}_t \rrbracket \, ds \quad \forall \mathbf{u}, \mathbf{v} \in \mathbf{V}_h, \end{aligned}$$

$$b(\mathbf{v}, q) := -(q, \operatorname{div} \mathbf{v})_\Omega \quad \forall \mathbf{v} \in \mathbf{V}_h, \quad \forall q \in \mathcal{Q}_h.$$

Consistency The consistency of the formulation (3.6) can be checked by means of the usual DG-machinery. In this case, it is sufficient to compare (3.1) and (3.7) and to observe that if

$(\mathbf{u}, p)$ is the solution of (1.5), then

$$A_h(\mathbf{u}, \mathbf{v}_h) \equiv a_h(\mathbf{u}, \mathbf{v}_h), \quad B_h(\mathbf{v}_h, p) \equiv b(\mathbf{v}_h, p), \quad \forall \mathbf{v}_h \in \mathbf{V}_h \subseteq \mathbf{H}_{0,n}(\text{div}; \Omega),$$

Further, it is evident that, $B_h(\mathbf{u}, q_h) \equiv b(\mathbf{u}, q_h)$ for all $q_h \in \mathcal{Q}_h$. Hence, as $(\mathbf{u}, p)$ verifies (3.2), it also verifies (3.6); that is,

$$(3.8) \quad \begin{cases} a_h(\mathbf{u}, \mathbf{v}) + b(\mathbf{v}, p) = (\mathbf{f}, \mathbf{v}) & \forall \mathbf{v} \in \mathbf{V}_h \\ b(\mathbf{u}, q) = 0 & \forall q \in \mathcal{Q}_h. \end{cases}$$

Thus, consistency is proved.

To prove the existence and uniqueness of the solution of (3.6) and to obtain the optimal error bounds, we need to define suitable norms. We define the following semi-norms

$$|\mathbf{v}|_{1,h}^2 = \sum_{T \in \mathcal{T}_h} \|\nabla \mathbf{v}\|_{0,T}^2, \quad \|\llbracket \mathbf{v} \rrbracket_*^2 := \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|\llbracket \mathbf{v} \rrbracket_{0,e}^2, \quad \forall \mathbf{v} \in \mathbf{H}^1(\mathcal{T}_h),$$

and norms

$$(3.9) \quad \begin{aligned} \|\mathbf{v}\|_{DG}^2 &:= 2\nu |\mathbf{v}|_{1,h}^2 + 2\nu \|\llbracket \mathbf{v}_t \rrbracket_*^2 & \mathbf{v} \in \mathbf{H}^1(\mathcal{T}_h), \\ \|\mathbf{v}\|^2 &:= \|\mathbf{v}\|_{DG}^2 + \sum_{T \in \mathcal{T}_h} 2\nu h_T^2 |\boldsymbol{\varepsilon}(\mathbf{v})|_{1,T}^2 & \mathbf{v} \in \mathbf{H}^2(\mathcal{T}_h). \end{aligned}$$

And, we remark that the seminorms defined in (3.9) are actually norms with the additional requirement that $\mathbf{v} \in \mathbf{H}_{0,n}(\text{div}; \Omega)$. We also observe that when restricted to discrete functions $\mathbf{v} \in \mathbf{V}_h$, the $\|\cdot\|_{DG}$ -norm and the $\|\cdot\|$ are equivalent (using inverse inequality). Continuity can easily be shown for both bilinear forms:

$$\begin{aligned} |a_h(\mathbf{u}, \mathbf{v})| &\leq \|\mathbf{u}\| \|\mathbf{v}\| & \forall \mathbf{u}, \mathbf{v} \in \mathbf{H}^2(\mathcal{T}_h), \\ |b(\mathbf{v}, q)| &\leq \|\mathbf{v}\|_{1,h} \|q\|_{0,\Omega} & \forall \mathbf{v} \in \mathbf{H}^1(\mathcal{T}_h), \quad q \in L^2(\Omega)/\mathbb{R}. \end{aligned}$$

Following [9], the existence and uniqueness of the approximate solution and optimal error bounds are guaranteed if the following two conditions are satisfied:

(H1): coercivity: $\exists \gamma > 0$ independent of the mesh size h such that

$$(3.10) \quad a_h(\mathbf{v}, \mathbf{v}) \geq \gamma \|\mathbf{v}\|_{DG}^2 \quad \forall \mathbf{v} \in \mathbf{V}_h.$$

(H2): inf-sup condition: $\exists \beta > 0$ independent of the mesh size h such that

$$(3.11) \quad \sup_{\mathbf{v} \in \mathbf{V}_h} \frac{(\text{div } \mathbf{v}, q_h)_\Omega}{\|\mathbf{v}\|_{DG}} \geq \beta \|q_h\|_{0,\Omega} \quad \forall q_h \in \mathcal{Q}_h.$$

Condition **(H2)** is a consequence of the *inf-sup* condition that holds for the continuous problem (1.5):

$$\exists \beta > 0 \text{ s.t. } \forall h, \forall q_h \in \mathcal{Q}_h \quad \exists \mathbf{v} \in \mathbf{H}^1(\Omega) : \text{div } \mathbf{v} = q_h \text{ and } \|\mathbf{v}\|_{1,\Omega} \leq \frac{1}{\beta} \|q_h\|_{0,\Omega}.$$

It is well known that for all the families considered here, an interpolant $\mathbf{v}^I \in \mathbf{V}_h$ of $\mathbf{v}$ exists that verifies (2.19) (in particular for $s = 1$), and

$$\operatorname{div} \mathbf{v}^I = \operatorname{div} \mathbf{v} (= q_h).$$

By observing that $[\![\mathbf{v}]\!] = 0$ on the internal edges as $\mathbf{v} \in \mathbf{H}^1(\Omega)$, and by using the Agmon trace inequality [2] and (2.19) (for $s = 1$), we have

$$(3.12) \quad |[\![\mathbf{v}^I]\!]_*|^2 := \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}_t^I]\!]\|_{0,e}^2 = \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}^I - \mathbf{v}]_t]\|_{0,e}^2 \leq C |\mathbf{v}|_{1,\Omega}^2.$$

Hence, again using (2.19), we deduce that

$$\|\mathbf{v}^I\|_{DG} \leq C |\mathbf{v}|_{1,\Omega}.$$

Thus (3.11) is proved.

To prove (3.10) we need a discrete Körn inequality, which we prove in the following lemma. Also see Appendix A for further comments on the validity of the result in three dimensions.

Lemma 3.2. *Let $\mathbf{V}_h$ be a piecewise polynomial subspace of $\mathbf{H}_{0,n}(\operatorname{div}; \Omega)$. Then, $\exists C_K > 0$ independent of h such that*

$$(3.13) \quad |\mathbf{v}|_{1,h}^2 \leq C_K \left(\|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0,\mathcal{T}_h}^2 + \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}_t]\!]\|_{0,e}^2 \right), \quad \forall \mathbf{v} \in \mathbf{V}_h.$$

Proof. To show (3.13), a direct application of [6, Inequality (1.12)] to $\mathbf{v} \in \mathbf{V}_h$ gives

$$(3.14) \quad |\mathbf{v}|_{1,h}^2 \leq C_K \left(\|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0,\mathcal{T}_h}^2 + \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}_t]\!]\|_{0,e}^2 + \sup_{\substack{\boldsymbol{\eta} \in \mathbf{RM}(\Omega) \\ \|\boldsymbol{\eta}\|_{0,\Gamma}=1, \int_\Gamma \boldsymbol{\eta} = 0}} \left(\int_\Gamma \mathbf{v} \cdot \boldsymbol{\eta} ds \right)^2 \right),$$

where $\mathbf{RM}(\Omega)$ is the space of rigid motions on Ω defined by

$$\mathbf{RM}(\Omega) = \{ \mathbf{a} + \mathbf{b}\mathbf{x} : \mathbf{a} \in \mathbb{R}^2 \quad \mathbf{b} \in so(2) \}$$

with $so(2)$ denoting the set of skew-symmetric 2×2 matrices.

We now show that the last term in (3.14) can be bounded by the first two. We claim that

$$(3.15) \quad \sup_{\substack{\boldsymbol{\eta} \in \mathbf{RM}(\Omega) \\ \|\boldsymbol{\eta}\|_{0,\Gamma}=1, \int_\Gamma \boldsymbol{\eta} = 0}} \left(\int_\Gamma \mathbf{v} \cdot \boldsymbol{\eta} ds \right)^2 \leq C \left(\|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0,\mathcal{T}_h}^2 + \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}_t]\!]\|_{0,e}^2 \right).$$

There are surely many ways of checking (3.15). Here, we propose one. For $\mathbf{v} \in \mathbf{V}_h$ and $\boldsymbol{\eta} \in \mathbf{RM}(\Omega)$ with $\int_\Gamma \boldsymbol{\eta} ds = 0$, we set

$$\mathcal{I}(\mathbf{v}, \boldsymbol{\eta}) := \int_\Gamma \mathbf{v} \cdot \boldsymbol{\eta} ds,$$

and we want to prove that

$$(3.16) \quad \mathcal{I}(\mathbf{v}, \boldsymbol{\eta}) \leq C \left(\|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0, \mathcal{T}_h}^2 + \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|[\![\mathbf{v}_t]\!] \|_{0, e}^2 \right)^{1/2} \|\boldsymbol{\eta}\|_{0, \Gamma}$$

will easily give (3.15) taking the supremum with respect to $\boldsymbol{\eta}$ with $\|\boldsymbol{\eta}\|_{0, \Gamma} = 1$. To prove (3.16) for every $\boldsymbol{\eta} \in \mathbf{RM}(\Omega)$ with $\int_{\Gamma} \boldsymbol{\eta} ds = 0$, we consider the following auxiliary elasticity problem:

$$(3.17) \quad \begin{cases} \boldsymbol{\tau} = \boldsymbol{\varepsilon}(\boldsymbol{\chi}) & \text{in } \Omega \\ \operatorname{div} \boldsymbol{\tau} = \mathbf{0} & \text{in } \Omega \\ \boldsymbol{\tau} \cdot \mathbf{n} = \boldsymbol{\eta} & \text{on } \Gamma. \end{cases}$$

The above problem has a unique solution $\boldsymbol{\chi}$ provided we complement problem (3.17) with the side condition

$$\int_{\Omega} \boldsymbol{\chi} \cdot \boldsymbol{\zeta} dx = 0 \quad \forall \boldsymbol{\zeta} \in \mathbf{RM}(\Omega).$$

Due to well-known results on the regularity of the solutions of PDE systems on polygons, the solution $\boldsymbol{\tau}$ of (3.17) (which, a priori, on a totally general domain would only be in $(L^2(\Omega))_{sym}^{2 \times 2}$) satisfies the following a priori estimate: *a $p > 2$ (depending on the geometry of Ω) and a constant C_p exist such that for all $\boldsymbol{\eta} \in \mathbf{RM}(\Omega)$ the corresponding $\boldsymbol{\tau}$ satisfies*

$$(3.18) \quad \|\boldsymbol{\tau}\|_{(L^p(\Omega))_{sym}^{2 \times 2}} \leq C_p \|\boldsymbol{\eta}\|_{0, \Gamma}.$$

The following technical proposition is the 2-dimensional version of Proposition A.1 given in Appendix A. The result stated here is, restricted to the case of interest, $d = 2$:

Proposition 3.3. *Let T be a triangle with minimum angle $\theta > 0$, and let e be an edge of T . Then for every $p > 2$ and for every integer k_{max} , a constant $C_{p, \theta, k_{max}}$ exists such that*

$$(3.19) \quad \int_e \mathbf{v} \cdot (\boldsymbol{\tau} \cdot \mathbf{n}) ds \leq C_{p, \theta, k_{max}} h_T^{-1/2} \|\mathbf{v}\|_{0, e} h_T^{\frac{p-2}{p}} \|\boldsymbol{\tau}\|_{0, p, T}$$

for every $\boldsymbol{\tau} \in (L^p(\Omega))_{sym}^{2 \times 2}$ having zero divergence and for every $\mathbf{v} \in \mathbf{P}^{k_{max}}(T)$.

Integration by parts together with the second equation of (3.17), (2.4) and (3.5), gives

$$(3.20) \quad \begin{aligned} \mathcal{I}(\mathbf{v}, \boldsymbol{\eta}) &= \int_{\Gamma} \mathbf{v} \cdot \boldsymbol{\eta} ds = \int_{\Gamma} \mathbf{v} \cdot (\boldsymbol{\tau} \cdot \mathbf{n}) ds \\ &= (\boldsymbol{\varepsilon}(\mathbf{v}) : \boldsymbol{\tau})_{\mathcal{T}_h} + (\mathbf{v}, \operatorname{div} \boldsymbol{\tau})_{\mathcal{T}_h} - \langle [\![\mathbf{v}_t]\!] : \{\boldsymbol{\tau}\} \rangle_{\mathcal{E}_h^o} \\ &= (\boldsymbol{\varepsilon}(\mathbf{v}) : \boldsymbol{\tau})_{\mathcal{T}_h} - \langle [\![\mathbf{v}_t]\!] : \{\boldsymbol{\tau}\} \rangle_{\mathcal{E}_h^o}. \end{aligned}$$

At this point, we can apply (3.19) to each e of the last term in (3.20) and then use the generalized Hölder inequality (with $q = 1/2$ and $r = 2p/(p-2)$, so that $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$).

Therefore, we obtain

$$(3.21) \quad \sum_{e \in \mathcal{E}_h^o} \int_e \llbracket \mathbf{v}_t \rrbracket : \{\boldsymbol{\tau}\} \, ds \leq C \left(\sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|\mathbf{v}_t\|_{0,e}^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}_h^o} \|\boldsymbol{\tau}\|_{0,p,T(e)}^p \right)^{1/p} \left(\sum_{e \in \mathcal{E}_h^o} h_e^{\frac{p-2}{p}r} \right)^{1/r} \\ \leq C \|\llbracket \mathbf{v}_t \rrbracket\|_* \|\boldsymbol{\tau}\|_{0,p,\Omega} \mu(\Omega)^{1/r}$$

where for each $e \in \mathcal{E}_h^o$ with $e = \partial T^+ \cap \partial T^-$, the set $T(e)$ refers to $T(e) := T^+ \cup T^-$. In the second line, $\mu(\Omega)$ denotes the measure of the domain Ω , whereas the constant C still depends on p , k_{max} and on the maximum angle in the decomposition $\mathcal{T}_h$.

From (3.20), (3.21), and the bound (3.18) we then obtain

$$(3.22) \quad |\mathcal{I}(\mathbf{v}, \boldsymbol{\eta})| \leq C (\|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0,\mathcal{T}_h} + \|\llbracket \mathbf{v}_t \rrbracket\|_*) \|\boldsymbol{\eta}\|_{0,\Gamma}$$

which gives (3.16). Thus the proof of the lemma is complete. $\square$

Remark 3.4. *The fact that in inequality (3.13) only the jumps over the interior edges $e \in \mathcal{E}_h^o$ (but not on the boundary edges) are included, prevents a direct and straightforward application of the results from [6]. The proof presented here is surely too elaborate, and we believe that a simpler proof is possible. However some of the machinery used here is likely to be of use elsewhere. Therefore, we decided that it would be worthwhile to present the proof we have obtained to date.*

The stability of $a_h(\cdot, \cdot)$ in the $\|\cdot\|_{DG}$ -norm can now be easily checked with the usual DG machinery. We have

$$\left| \int_e \{\boldsymbol{\varepsilon}(\mathbf{v})\} : \llbracket \mathbf{v}_t \rrbracket \, ds \right| \leq h^{1/2} \|\{\boldsymbol{\varepsilon}(\mathbf{v})\}\|_{0,e} h^{-1/2} \|\llbracket \mathbf{v}_t \rrbracket\|_{0,e},$$

which when we proceed as in [3] (or as in (3.21) with $p = 2$) yields

$$(3.23) \quad \left| \sum_{e \in \mathcal{E}_h^o} \int_e \{\boldsymbol{\varepsilon}(\mathbf{v})\} : \llbracket \mathbf{v}_t \rrbracket \, ds \right| \leq C |\mathbf{v}|_{1,h} \|\llbracket \mathbf{v}_t \rrbracket\|_*.$$

Using (3.23) in (3.7), we then have

$$a_h(\mathbf{v}, \mathbf{v}) \geq 2\nu \|\boldsymbol{\varepsilon}(\mathbf{v})\|_{0,\mathcal{T}_h}^2 + 2\nu \alpha \|\llbracket \mathbf{v}_t \rrbracket\|_*^2 - 4\nu C |\mathbf{v}|_{1,\mathcal{T}_h} \|\llbracket \mathbf{v}_t \rrbracket\|_*.$$

Now using the Körn inequality (3.13) and the usual arithmetic-geometric mean inequality, we easily have a big enough α :

$$a_h(\mathbf{v}, \mathbf{v}) \geq \gamma \|\mathbf{v}\|_{DG}^2 \quad \forall \mathbf{v} \in \mathbf{V}_h.$$

We close this section with the following theorem.

Theorem 3.5. *Let $(\mathbf{V}_h, \mathcal{Q}_h)$ be as in one of our three examples. Then problem (3.6) has a unique solution $(\mathbf{u}_h, p_h) \in \mathbf{V}_h \times \mathcal{Q}_h$ that verifies*

$$(3.24) \quad \operatorname{div} \mathbf{u}_h = 0 \quad \text{in } \Omega.$$

Moreover, there exists a positive constant C , independent of h , such that for every $\mathbf{v}_h \in \mathbf{V}_h$ with $\operatorname{div} \mathbf{v}_h = 0$ and for every $q_h \in \mathcal{Q}_h$ the following estimate holds:

$$(3.25) \quad \|\mathbf{u} - \mathbf{u}_h\|_{DG} \leq C \|\mathbf{u} - \mathbf{v}_h\|_{DG}, \quad \|p - p_h\|_{0,\Omega} \leq C (\|p - q_h\|_{0,\Omega} + \|\mathbf{u} - \mathbf{v}_h\|_{DG}),$$

with $(\mathbf{u}, p)$ solution of (1.5).

Proof. The existence and uniqueness of the solution of (3.6) follow from (3.10)-(3.11). The divergence-free property (3.24) is implied by (2.14), which holds for all our choices of spaces. Let $\mathbf{v}_h \in \mathbf{V}_h$ also be divergence-free; then we obviously have that $b(\mathbf{v}_h - \mathbf{u}_h, q) = 0$ for every $q \in L^2(\Omega)/\mathbb{R}$. In particular, $b(\mathbf{v}_h - \mathbf{u}_h, p - p_h) = 0$. Hence, from the coercivity (3.10), consistency (3.8), and continuity of $a_h(\cdot, \cdot)$ we deduce immediately

$$\gamma \|\mathbf{v}_h - \mathbf{u}_h\|_{DG}^2 \leq a_h(\mathbf{v}_h - \mathbf{u}_h, \mathbf{v}_h - \mathbf{u}_h) = a_h(\mathbf{v}_h - \mathbf{u}, \mathbf{v}_h - \mathbf{u}_h) \leq \|\mathbf{v}_h - \mathbf{u}\|_{DG} \|\mathbf{v}_h - \mathbf{u}_h\|_{DG}.$$

On the same basis we deduce that the first estimate in (3.25) follows by triangle inequality. For every $\mathbf{w}_h \in \mathbf{V}_h$, using the consistency and continuity of $a_h(\cdot, \cdot)$, we have

$$(3.26) \quad \begin{aligned} b(\mathbf{w}_h, q_h - p_h) &= b(\mathbf{w}_h, q_h - p) + b(\mathbf{w}_h, p - p_h) = b(\mathbf{w}_h, q_h - p) - a_h(\mathbf{u} - \mathbf{u}_h, \mathbf{w}_h) \\ &\leq (\|q_h - p\|_{0,\Omega} + \|\mathbf{u} - \mathbf{u}_h\|_{DG}) \|\mathbf{w}_h\|_{DG}. \end{aligned}$$

By dividing (3.26) by $\|\mathbf{w}_h\|_{DG}$ and then using the *inf-sup* condition (3.11), we immediately deduce that

$$\beta \|q_h - p_h\|_{0,\Omega} \leq \|q_h - p\|_{0,\Omega} + \|\mathbf{u} - \mathbf{u}_h\|_{DG},$$

and that the second estimate in (3.25) follows again by triangle inequality. $\square$

Remark 3.6. In the assumptions of Theorem 3.5, we could obviously consider any trio of finite element spaces satisfying **H0**. However, for choices like **RT**₀, not considered in our three examples, the estimate (3.25) could be meaningless, as the term $\|\mathbf{u} - \mathbf{v}_h\|_{DG}$ does not, in general, go to zero with h . Still, this choice could be profitably used, in some cases, as a preconditioner, as it does satisfy **H0**, **H1**, and **H2**.

4. DISCRETE HELMHOLTZ DECOMPOSITIONS

So far, we have assumed that the computational domain Ω is a polygon. From now on, for the sake of simplicity, we are going to work under the stronger assumption that Ω is a *convex polygon*. As is well known, this allows the use of better regularity results, and in particular the H^2 -regularity for elliptic second-order operators.

Following [9] we define the discrete gradient operator $\mathcal{G}_h : \mathcal{Q}_h \rightarrow \mathbf{V}_h$ as

$$(4.1) \quad (\mathcal{G}_h q_h, \mathbf{v}_h)_{0,\Omega} := -(q_h, \operatorname{div} \mathbf{v}_h)_{0,\Omega} \quad \forall \mathbf{v}_h \in \mathbf{V}_h.$$

Lemma 4.1. Assume that together the three spaces $(\mathbf{V}_h, \mathcal{Q}_h, \mathcal{N}_h)$ satisfy assumption **H0** (given in Definition 3.1). Then for any $\mathbf{v}_h \in \mathbf{V}_h$ a unique $q_h \in \mathcal{Q}_h$ and a unique $\varphi_h \in \mathcal{N}_h$ exist such that

$$(4.2) \quad \mathbf{v}_h = \mathcal{G}_h q_h + \operatorname{curl} \varphi_h,$$

that is,

$$(4.3) \quad \mathbf{V}_h = \mathcal{G}_h(\mathcal{Q}_h) \oplus \mathbf{curl} \mathcal{N}_h.$$

Moreover, a constant C independent of h exists, such that the following estimate holds:

$$(4.4) \quad \|\mathcal{G}_h q_h\|_{DG} \leq C \|\operatorname{div} \mathbf{v}_h\|_{0,\Omega}.$$

Proof. For $\mathbf{v}_h \in \mathbf{V}_h$, consider the auxiliary problem:

$$(4.5) \quad -\Delta q = \operatorname{div} \mathbf{v}_h \quad \text{in } \Omega, \quad \frac{\partial q}{\partial n} = 0 \quad \text{on } \partial\Omega, \quad \text{and} \quad \int_{\Omega} q \, dx = 0.$$

Owing to the boundary conditions in $\mathbf{V}_h$, we have that $\operatorname{div} \mathbf{v}_h$ has zero mean value in Ω . Hence, problem (4.5) has a unique solution, that satisfies

$$(4.6) \quad \|q\|_{2,\Omega} \leq C_{reg} \|\operatorname{div} \mathbf{v}_h\|_{0,\Omega}.$$

We write (4.5) in mixed form:

$$\boldsymbol{\sigma} = -\nabla q \text{ in } \Omega, \quad \operatorname{div} \boldsymbol{\sigma} = \operatorname{div} \mathbf{v}_h \text{ in } \Omega, \quad \boldsymbol{\sigma} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega.$$

and, we consider directly the approximation of the mixed formulation: Find $(\boldsymbol{\sigma}_h, q_h) \in \mathbf{V}_h \times \mathcal{Q}_h$ such that :

$$(4.7) \quad \begin{cases} (\boldsymbol{\sigma}_h, \boldsymbol{\tau})_{0,\Omega} - (q_h, \operatorname{div} \boldsymbol{\tau})_{0,\Omega} = 0 & \forall \boldsymbol{\tau} \in \mathbf{V}_h, \\ (\operatorname{div} \boldsymbol{\sigma}_h, s_h)_{0,\Omega} = (\operatorname{div} \mathbf{v}_h, s_h)_{0,\Omega} & \forall s_h \in \mathcal{Q}_h. \end{cases}$$

Problem (4.7) obviously has a unique solution, which moreover satisfies

$$(4.8) \quad \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_h\|_{0,\Omega} \leq C h \|\boldsymbol{\sigma}\|_{1,\Omega} \leq C C_{reg} h \|\operatorname{div} \mathbf{v}_h\|_{0,\Omega},$$

given that (4.6) was used in the last step. As both $\mathbf{v}_h$ and $\boldsymbol{\sigma}_h$ are in $\mathbf{V}_h$ (and as (2.14) holds,) the second equation in (4.7) directly implies that

$$\operatorname{div} (\boldsymbol{\sigma}_h - \mathbf{v}_h) = 0.$$

Hence, the exact sequence (2.8) implies that

$$(4.9) \quad \text{a unique } \varphi_h \in \mathcal{N}_h \text{ exists such that } \boldsymbol{\sigma}_h - \mathbf{v}_h = \mathbf{curl} \varphi_h.$$

Next, by using the first equation in (4.7) and then applying definition (4.1), we deduce that

$$(\boldsymbol{\sigma}_h, \boldsymbol{\tau})_{0,\Omega} = (q_h, \operatorname{div} \boldsymbol{\tau})_{0,\Omega} = -(\mathcal{G}_h q_h, \boldsymbol{\tau})_{0,\Omega} \quad \forall \boldsymbol{\tau} \in \mathbf{V}_h,$$

which implies $\boldsymbol{\sigma}_h = -\mathcal{G}_h q_h$, that joined to (4.9) gives (4.2).

In order to prove (4.4), we recall that

$$(4.10) \quad \|\mathcal{G}_h q_h\|_{DG}^2 = \|\boldsymbol{\sigma}_h\|_{DG}^2 = \|\nabla \boldsymbol{\sigma}_h\|_{0,\mathcal{T}_h}^2 + \|[(\boldsymbol{\sigma}_h)_t]\|_*^2.$$

For the first term, by adding and subtracting the interpolant σ^I of σ and then using inverse inequality and (2.19), we have:

$$(4.11) \quad \begin{aligned} \|\nabla \sigma_h\|_{0,\mathcal{T}_h} &\leq \|\nabla(\sigma_h - \sigma^I)\|_{0,\mathcal{T}_h} + \|\nabla \sigma^I\|_{0,\mathcal{T}_h} \\ &\leq C_{inv} h^{-1} \|\sigma_h - \sigma^I\|_{0,\mathcal{T}_h} + C \|\nabla \sigma\|_{0,\mathcal{T}_h}. \end{aligned}$$

From triangle inequality, (4.8), and standard approximation properties (see (2.19)), we have

$$(4.12) \quad \|\nabla \sigma_h\|_{0,\mathcal{T}_h} \leq C \|\operatorname{div} \mathbf{v}_h\|_{0,\Omega}.$$

The jump term in (4.10) is estimated similarly. First, we remark that $\sigma = -\nabla q$ with $q \in H^2(\Omega)$ so that $[\![\sigma]\!] = 0$ on each $e \in \mathcal{E}_h^o$, therefore

$$|[\![(\sigma_h)_t]\!]_*|^2 = |[\![(\sigma_h)_t - \sigma_t]\!]_*|^2.$$

Then, using Agmon trace inequalities (4.8) and the boundedness of σ_h and σ , we have

$$\begin{aligned} |[\![(\sigma_h)_t - \sigma_t]\!]_*|^2 &= \sum_{e \in \mathcal{E}_h^o} h_e^{-1} |[\![(\sigma_h)_t - \sigma_t]\!]_{0,e}|^2 \\ &\leq C_t h^{-2} \|\sigma_h - \sigma\|_{0,\mathcal{T}_h}^2 + C_t \|\nabla(\sigma_h - \sigma)\|_{0,\mathcal{T}_h}^2 \\ &\leq C C_{reg} \|\operatorname{div} \mathbf{v}_h\|_{0,\Omega}^2. \end{aligned}$$

Thus the proof is complete. $\square$

5. PRECONDITIONER: FICTITIOUS SPACE LEMMA AND AUXILIARY SPACE FRAMEWORK

5.1. Preconditioner for the semi-definite system. Assume V is a Hilbert space equipped with the norm $\|\cdot\|_V$ and that $A : V \mapsto V'$ is a bounded linear operator. We define the bilinear form

$$(u, v)_A = \langle Au, v \rangle.$$

We say A is symmetric if the bilinear form $(u, v)_A$ is symmetric. We say that A is semi-positive definite if

$$(v, v)_A \geq 0, \quad \forall v \in V$$

and $\alpha > 0$ exists such that

$$(v, v)_A \geq \alpha \|v\|_{V/N(A)}^2, \quad \forall v \in V/N(A).$$

And, A is SPD if $\alpha > 0$ exists such that

$$(v, v)_A \geq \alpha \|v\|_V^2, \quad \forall v \in V.$$

One useful property of the symmetric a semi-positive definite operator is that

$$(5.1) \quad Av = 0 \text{ iff } \langle Av, v \rangle = 0.$$

A preconditioner for A is another symmetric semi-positive definite operator $B : V' \mapsto V$. Again, we consider the bilinear form

$$(f, g)_B = \langle f, Bg \rangle.$$

The operator $BA : V \mapsto V$ satisfies

$$(BAu, v)_A = \langle Av, BAu \rangle = (Au, Av)_B.$$

Lemma 5.1. *If $A : V \mapsto V'$ and $B : V' \mapsto V$ are both symmetric semi-positive definite such that B is positive definite on $R(A)$, then*

- (1) $B : R(A) \mapsto R(BA)$ is an isomorphism (with the inverse satisfying trivially that $B^{-1}(BAv) = Av$).
- (2) The bilinear form $(\cdot, \cdot)_{B^{-1}}$ defines an inner product on $R(BA)$.
- (3) The bilinear form $(\cdot, \cdot)_A$ defines an inner product on $R(BA)$.
- (4) BA is symmetric positive definite on $R(BA)$ with either of the above two inner products.

Proof. All these results are pretty obvious, and their proofs are similar. Let us give the proof for 3 as an example.

We only need to verify that $(\cdot, \cdot)_A$ is positive definite on $R(BA)$. If $v \in R(BA)$ is such that $(v, v)_A = 0$, then, by (5.1), we have $Av = 0$. We write $v = BA w$ for some $w \in V$, then $ABAw = 0$ and hence $(Aw, Aw)_B = 0$. As B is positive definite on $R(A)$, we have $Aw = 0$. Thus, $v = ABAw = 0$, as desired. $\square$

For the system $Au = f$, we can apply the preconditioner B and the preconditioned conjugate gradient (PCG) method with respect to the inner product $(\cdot, \cdot)_{B^{-1}}$ with the following convergence estimate:

$$\|u - u^k\|_A \leq 2 \left(\frac{\sqrt{\kappa(BA)} - 1}{\sqrt{\kappa(BA)} + 1} \right)^k \|u - u^0\|_A.$$

The condition number can then be estimated by $\kappa(BA) \leq c_1/c_0$, either where

$$c_0(v, v)_{B^{-1}} \leq (BAv, v)_{B^{-1}} \leq c_1(v, v)_{B^{-1}}, \quad \forall v \in R(BA),$$

or equivalently where

$$c_0(w, w)_B \leq (Bw, Bw)_A \leq c_1(w, w)_B, \quad \forall w \in R(A),$$

or where

$$c_1^{-1}(v, v)_A \leq (B^{-1}v, v) \leq c_0^{-1}(v, v)_A \quad \forall v \in R(BA).$$

5.2. Fictitious space lemma and generalizations. Let us present and prove a refined version of the Fictitious Space Lemma originally proposed by Nepomnyaschikh [16] (see also [20]).

Lemma 5.2. *Let $\tilde{V}$ and V be two Hilbert spaces, and let $\Pi : \tilde{V} \mapsto V$ be a surjective map. Let $\tilde{B} : \tilde{V}' \mapsto \tilde{V}$ be a symmetric and positive definite operator. Then $B := \Pi \tilde{B} \Pi'$ is also*

symmetric and positive definite (here $\Pi' : V' \mapsto \tilde{V}'$ is such that $\langle \Pi'g, \tilde{v} \rangle = \langle g, \Pi\tilde{v} \rangle$, $g \in V'$ and $\tilde{v} \in \tilde{V}$). Furthermore,

$$\langle B^{-1}v, v \rangle = \inf_{\Pi\tilde{v}=v} \langle \tilde{B}^{-1}\tilde{v}, \tilde{v} \rangle.$$

Proof. It is obvious that B is symmetric and positive semi-definite. Note that if $v \in V$ such that $\langle Bv, v \rangle = 0$, then $\langle \tilde{B}\Pi'v, \Pi'v \rangle = \langle Bv, v \rangle = 0$. This means that $\Pi'v = 0$ as $\tilde{B}$ is SPD. Hence, $v = 0$ as Π' is injective. This proves that B is positive definite.

For any $\tilde{v} \in \tilde{V}$, let $v = \Pi\tilde{v}$ and $\tilde{v}^* = \tilde{B}\Pi'B^{-1}v$. As we obviously have $\Pi\tilde{v}^* = v$, we can write $\tilde{v} = \tilde{v}^* + \tilde{w}$ with $\Pi\tilde{w} = 0$. Thus,

$$\begin{aligned} \inf_{\Pi\tilde{v}=v} \langle \tilde{B}^{-1}\tilde{v}, \tilde{v} \rangle &= \inf_{\Pi\tilde{w}=0} \langle \tilde{B}^{-1}(\tilde{v}^* + \tilde{w}), \tilde{v}^* + \tilde{w} \rangle \\ &= \langle \tilde{B}^{-1}\tilde{v}^*, \tilde{v}^* \rangle + \inf_{\Pi\tilde{w}=0} \left(\langle \tilde{B}^{-1}\tilde{w}, \tilde{w} \rangle + 2\langle \tilde{B}^{-1}\tilde{v}^*, \tilde{w} \rangle \right) \end{aligned}$$

From the definition of $\tilde{v}^*$ we have

$$\langle \tilde{B}^{-1}\tilde{v}^*, \tilde{v}^* \rangle = \langle B^{-1}v, \Pi\tilde{v}^* \rangle = \langle B^{-1}v, v \rangle,$$

and also

$$\langle \tilde{B}^{-1}\tilde{v}^*, \tilde{w} \rangle = \langle \tilde{B}^{-1}\tilde{B}\Pi'B^{-1}v, \tilde{w} \rangle = \langle \Pi'B^{-1}v, \tilde{w} \rangle = \langle B^{-1}v, \Pi\tilde{w} \rangle = 0.$$

The last two identities lead to the desired result. $\square$

Theorem 5.3. Assume that $\tilde{A} : \tilde{V} \mapsto \tilde{V}'$ and $A : V \mapsto V'$ are symmetric semi-definite operators. We assume that $\Pi : \tilde{V} \mapsto V$ is surjective and that $\Pi(N(\tilde{A})) = N(A)$. Then for any SPD operator $\tilde{B} : \tilde{V}' \mapsto \tilde{V}$, we have, for $B = \Pi\tilde{B}\Pi'$,

$$\kappa(BA) \leq \kappa(\Pi)\kappa(\tilde{B}\tilde{A}).$$

Here $\kappa(\Pi)$ is the smallest ratio c_1/c_0 that satisfies

$$(5.2) \quad c_1^{-1}\langle Av, v \rangle \leq \inf_{\Pi\tilde{v}=v} \langle \tilde{A}\tilde{v}, \tilde{v} \rangle \leq c_0^{-1}\langle Av, v \rangle, \quad \forall v \in R(BA).$$

Proof. Denote $\kappa(\tilde{B}\tilde{A}) = b_1/b_0$ with b_1 and b_0 satisfying

$$b_1^{-1}(\tilde{v}, \tilde{v})_{\tilde{A}} \leq (\tilde{B}^{-1}\tilde{v}, \tilde{v}) \leq b_0^{-1}(\tilde{v}, \tilde{v})_{\tilde{A}}, \quad \forall \tilde{v} \in R(\tilde{B}\tilde{A}).$$

By (5.2), we obtain

$$b_1^{-1}c_1^{-1}\|v\|_A^2 \leq \inf_{\Pi\tilde{v}=v, \tilde{v} \in R(\tilde{B}\tilde{A})} (\tilde{B}^{-1}\tilde{v}, \tilde{v}) \leq b_0^{-1}c_0^{-1}\|v\|_A^2, \quad \forall v \in R(BA).$$

By the assumption that $\Pi(N(\tilde{A})) = N(A)$, we can prove that $\Pi'(R(A)) \subset R(\tilde{A})$ and

$$\{\tilde{v} | \Pi\tilde{v} = v \in R(BA)\} = \{\tilde{v} | \Pi\tilde{v} = v \in R(BA), \tilde{v} \in R(\tilde{B}\tilde{A})\}.$$

By Lemma 5.2,

$$\inf_{\Pi\tilde{v}=v, \tilde{v} \in R(\tilde{B}\tilde{A})} (\tilde{B}^{-1}\tilde{v}, \tilde{v}) = \inf_{\Pi\tilde{v}=v} (\tilde{B}^{-1}\tilde{v}, \tilde{v}) = (B^{-1}v, v), \quad \forall v \in R(BA).$$

Therefore,

$$b_1^{-1}c_1^{-1}\|v\|_A^2 \leq (B^{-1}v, v) \leq b_0^{-1}c_0^{-1}\|v\|_A^2 \quad \forall v \in R(BA).$$

□

Theorem 5.4. *Assume that the following two conditions are satisfied for Π . First,*

$$\|\Pi\tilde{v}\|_A \leq c_1\|\tilde{v}\|_{\tilde{A}}, \quad \forall \tilde{v} \in \tilde{V}.$$

Second, for any $v \in V$, $\tilde{v} \in \tilde{V}$ exists such that $\Pi\tilde{v} = v$ and

$$\|\tilde{v}\|_{\tilde{A}} \leq c_0\|v\|_A.$$

Then $\kappa(\Pi) \leq c_1/c_0$ and, under the assumptions of Theorem 5.3,

$$\kappa(BA) \leq \left(\frac{c_1}{c_0}\right)^2 \kappa(\tilde{B}\tilde{A}).$$

Remark 5.5. *In view of the application of the above results to our case (as we shall see in the next subsection), it would have been enough to restrict ourselves to the symmetric positive definite case (instead of the semi-definite case treated in the last two subsections). However we preferred to have them in the present more general setting, as in this form they are likely to be useful in many other circumstances (starting, as natural, from the extension of the present theory to the three-dimensional case).*

5.3. Application to our problem. In this section we design a simple preconditioner for the linear system resulting from the approximation of the Stokes problem (1.5) defined in (3.6)-(3.7). Note that the bilinear form $a_h(\cdot, \cdot)$ defined in (3.7) provides a discretization of the vector Laplacian problem

$$-\mathbf{div}(2\nu\boldsymbol{\varepsilon}(\mathbf{u})) = \mathbf{f} \quad \text{in } \Omega, \quad \mathbf{u} \cdot \mathbf{n} = 0, \quad (\boldsymbol{\varepsilon}(\mathbf{u}) \cdot \mathbf{n}) \cdot \mathbf{t} = 0 \quad \text{on } \Gamma.$$

We denote by A the operator associated with $a_h(\cdot, \cdot)$. As the solution $\mathbf{u}_h \in \mathbf{V}_h$ of (3.6) is divergence-free, the discrete Helmholtz decomposition (4.2) implies that

$$\text{a unique } \psi_h \in \mathcal{N}_h \quad \text{exists such that } \mathbf{u}_h = \mathbf{curl} \psi_h.$$

At this point, it is convenient to introduce the space $\overset{o}{\mathbf{V}}_h$ as

$$(5.3) \quad \overset{o}{\mathbf{V}}_h := \mathbf{V}_h \cap \mathbf{H}_0(\text{div}^0; \Omega).$$

And, we note that as the sequence (2.8) is exact, we have

$$(5.4) \quad \overset{o}{\mathbf{V}}_h \equiv \mathbf{curl} \mathcal{N}_h,$$

and that the mapping is one-to-one. Therefore, restricting the bilinear form $a_h(\cdot, \cdot)$ to $\overset{o}{\mathbf{V}}_h$, in the spirit of Remark 2.1, corresponds here to restricting the trial and test space to $\overset{o}{\mathbf{V}}_h \equiv$

$\mathbf{curl}(\mathcal{N}_h)$. The discrete problem (3.6) then reduces to the following problem: *Find $\psi_h \in \overset{o}{V}_h$ such that*

$$(5.5) \quad a_h(\psi_h, \varphi_h) = (\mathbf{f}, \varphi_h) \quad \forall \varphi_h \in \overset{o}{V}_h$$

Defining the operator $A_h : \overset{o}{V}_h \mapsto \overset{o}{V}_h'$ by $\langle A_h \psi_h, \varphi_h \rangle = a_h(\psi_h, \varphi_h)$, $\psi_h, \varphi_h \in \overset{o}{V}_h$, we can write (5.5) as

$$A_h \psi_h = f_h.$$

We now use the original space $\mathbf{V}_h$ as the auxiliary space for $\overset{o}{V}_h$. Define $\tilde{A}_h : \mathbf{V}_h \mapsto \mathbf{V}_h'$ by $\langle \tilde{A}_h u_h, v_h \rangle = a_h(u_h, v_h)$, $u_h, v_h \in \mathbf{V}_h$. We note that $\tilde{A}_h$ is a discrete Laplacian. We assume that $\tilde{B}_h$ is an optimal preconditioner for $\tilde{A}_h$.

We now define the operator

$$(5.6) \quad \Pi_h : \mathbf{V}_h \longrightarrow \overset{o}{V}_h \equiv \mathbf{curl}(\mathcal{N}_h)$$

which maps each $\mathbf{v}_h \in \mathbf{V}_h$ to its divergence free part in $\overset{o}{V}_h$, according to (4.2). Note that Π_h is a surjective operator and that Π_h acts as the identity on the subspace $\overset{o}{V}_h$. The auxiliary space preconditioner for A_h is then defined by

$$(5.7) \quad B_h = \Pi_h \tilde{B}_h \Pi_h^*.$$

Lemma 5.6. *Assume that the spaces $(\mathbf{V}_h, \mathcal{Q}_h, \mathcal{N}_h)$ satisfy assumption **H0**. Then B_h given by (5.7) is an optimal preconditioner for A_h as long as $\tilde{B}_h$ is an optimal preconditioner for $\tilde{A}_h$.*

Proof. Following the auxiliary space techniques (Theorem 5.4), we need to check that the following two properties are satisfied:

(A1): Local Stability: there exists a positive constant C_1 independent of h such that

$$(5.8) \quad \|\Pi_h \mathbf{v}_h\|_{DG} \leq C_1 \|\mathbf{v}_h\|_{DG} \quad \forall \mathbf{v}_h \in \mathbf{V}_h$$

(A2): Stable decomposition: there exists a positive constant C_2 independent of h such

that for any $\mathbf{w}_h \in \overset{o}{V}_h$ there exists $\mathbf{v}_h \in \mathbf{V}_h$ such that $\Pi_h \mathbf{v}_h = \mathbf{w}_h$ and

$$(5.9) \quad \|\mathbf{v}_h\|_{DG} \leq C_2 \|\mathbf{w}_h\|_{DG}.$$

To prove (5.8) from the Helmholtz decomposition (4.2) and the definition (5.6) of Π_h , we have

$$(5.10) \quad \mathbf{v}_h = \mathcal{G}_h q_h + \mathbf{curl} \varphi_h = \mathcal{G}_h q_h + \Pi_h \mathbf{v}_h.$$

Using estimate (4.4) from Lemma 4.1 and the clear fact that $\text{div} \mathbf{v}_h$ is the trace of $\boldsymbol{\varepsilon}(\mathbf{v}_h)$, we have

$$(5.11) \quad \|\mathcal{G}_h q_h\|_{DG} \leq C \|\text{div} \mathbf{v}_h\|_{0,\Omega} \leq C \|\boldsymbol{\varepsilon}(\mathbf{v}_h)\|_{0,\mathcal{T}_h} \leq C \|\mathbf{v}_h\|_{DG}.$$

Hence, (5.8) follows from (5.10) and (5.11):

$$\|\Pi_h \mathbf{v}_h\|_{DG} = \|\mathbf{v}_h - \mathcal{G}_h q_h\|_{DG} \leq \|\mathbf{v}_h\|_{DG} + \|\mathcal{G}_h q_h\|_{DG} \leq C \|\mathbf{v}_h\|_{DG}.$$

Finally, the inequality (5.9) holds with $C_2 = 1$ by taking $\mathbf{v}_h = \mathbf{w}_h$. $\square$

6. NUMERICAL EXPERIMENTS

6.1. Setup. The tests presented in this section use discretization by the lowest order, namely, $\mathbf{BDM}_1$ elements paired with piece-wise constant space for the pressure. They verify the *a priori* estimates given in Theorem 3.5 and confirm the uniform bound on the condition number of the preconditioned system for the velocity.

As previously set up, the discrete problem under consideration is given by equation (3.6) with bilinear forms $a_h(\cdot, \cdot)$ and $b(\cdot, \cdot)$ defined in (3.7). In the numerical tests presented here, we take $\nu = 1/2$ and the penalty parameter $\alpha = 4$ in (3.7). We present two sets of tests




Figure 6.1. Meshes used in the tests for the unit square domain $\Omega = (0, 1) \times (0, 1)$

with A corresponding to the Stokes equation discretized on a sequence of successively refined unstructured meshes as shown in Figures 6.1–6.2. On the square the coarsest mesh (level of refinement $J = 0$) has 160 elements and 97 vertices with 448 BDM degrees of freedom. The finer triangulations of the square domain are obtained via $1, \dots, 5$ regular refinements (every element divided in 4) and the finest one is with 163,840 elements, 82,433 vertices and 490,496 BDM degrees of freedom. Similarly for the L -shaped domain we start with a coarsest grid ($J = 0$) with 64 vertices and 97 elements. For the L -shaped domain the finest grid (for $J = 5$) has 99,328 elements, 50,129 vertices and 297,056 $\mathbf{BDM}_1$ degrees of freedom. In the computations, we approximate the velocity component $\mathbf{u}_h$ of the solution of the Stokes equation by solving several simpler equations (such as scalar Laplace equations). After we obtain the velocity, the pressure then is found via a postprocessing step at low




Figure 6.2. Meshes used in the tests for the L -shaped domain $\Omega = ((0, 1) \times (0, 1)) \setminus ([\frac{1}{2}, 1) \times [\frac{1}{2}, 1))$

computational cost. Further, for this sequence of grids the $\mathbf{BDM}_1$ interpolant of a function $\mathbf{v}$ on the k -th grid is denoted by $\mathbf{v}^{I_k}$. Accordingly the piece-wise constant, L_2 -orthogonal projection of p is denoted by p^{I_k} . We also use the notation $(\mathbf{u}_k, p_k)$ for the solution of (3.6) on the k -th grid, $k = 0, \dots, 5$.

6.2. Discretization error. We now present several tests related to the error estimates given in the previous sections. We computed and tabulated approximations of the order of convergence of the discrete solution in different norms. These approximations are denoted by $\gamma_0 \approx \beta_0$, $\gamma_{DG} \approx \beta_{DG}$, $\gamma_p \approx \beta_p$, and $\gamma_* \approx \beta_*$. The actual orders of convergence β_0 , β_{DG} , β_p , and β_* are

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_{0,\Omega} &\approx C(\mathbf{u})h^{\beta_0}, & \|\mathbf{u} - \mathbf{u}_h\|_{DG} &\approx C(\mathbf{u})h^{\beta_{DG}}, \\ \|p - p_h\|_{0,\Omega} &\approx C(\mathbf{u}, p)h^{\beta_p}, & \|[\![\mathbf{u}_h]\!]\|_* &\approx C(\mathbf{u})h^{\beta_*}. \end{aligned}$$

Here, as in (3.12), we denote

$$|[\![\mathbf{v}]\!]|_*^2 = \sum_{e \in \mathcal{E}_h^o} h_e^{-1} \int_e [\![\mathbf{u}_t]\!]^2 ds.$$

Note that β_* is the order with which the jumps in the approximate solution (not in the error) go to zero.

We present two sets of experiments to illustrate the results given in Theorem 3.5. First, we consider the exact given solution and calculate the right-hand side and the boundary conditions from this solution. We set

$$(6.1) \quad \phi = xy(1-x)(2x-1)(y-1)(2y-1), \quad \mathbf{u} = \mathbf{curl}\phi.$$

Square domain						<i>L</i> -shaped domain					
k	1	2	3	4	5	k	1	2	3	4	5
γ_0	1.56	1.79	1.91	1.96	1.98	γ_0	1.43	1.64	1.84	1.93	1.97
γ_{DG}	1.32	1.58	1.66	1.66	1.62	γ_{DG}	1.17	1.47	1.60	1.62	1.60
γ_p	0.43	0.70	0.86	0.94	0.97	γ_p	0.24	0.52	0.79	0.91	0.96
γ_*	0.70	0.85	0.94	0.97	0.99	γ_*	0.64	0.76	0.89	0.95	0.98

Table 6.1. Approximate order of convergence for the difference $(\mathbf{u}^I - \mathbf{u}_h)$ and $(p^I - p_h)$ and the jumps $|\llbracket \mathbf{u}_h \rrbracket|_*$ for the square and *L*-shaped domains. Here, $\mathbf{u}$ and p are given in (6.1) and (6.2).

Clearly, the function ϕ vanishes on the boundary of both the domains under consideration and we take $\mathbf{u}$ defined in (6.1) as exact solution for the velocity for both the square and the *L*-shaped domains. For the pressure we choose as exact solutions functions with zero mean value and select p different for the square and the *L*-shaped domain, namely

$$(6.2) \quad \begin{aligned} p &= x^2 - 3y^2 + \frac{8}{3}xy, & (\text{square domain}), \\ p &= x^2 - 3y^2 + \frac{24}{7}xy, & (L\text{-shaped domain}). \end{aligned}$$

The right hand side $\mathbf{f}$ is calculated by plugging $(\mathbf{u}, p)$ defined in (6.1)–(6.2) in (1.1). Table 6.1 shows tabulation of the order of convergence of $(\mathbf{u}_h, p_h)$ to $(\mathbf{u}^I, p^I)$ for both the square domain and the *L*-shaped domain. The values approximating the order of convergence displayed in Table 6.1 are

$$\begin{aligned} \gamma &= \log_2 \frac{\|\mathbf{u}_{k-1}^I - \mathbf{u}_{k-1}\|}{\|\mathbf{u}_k^I - \mathbf{u}_k\|}, & \gamma_* &= \log_2 \frac{|\llbracket \mathbf{u}_k \rrbracket|_*}{|\llbracket \mathbf{u}_{k-1} \rrbracket|_*}, \\ \gamma_p &= \log_2 \frac{\|p_{k-1}^I - p_{k-1}\|_{0,\Omega}}{\|p_k^I - p_k\|_{0,\Omega}}, & k &= 1, \dots, 5. \end{aligned}$$

Here $\|\cdot\|$ stands for any of the *DG* or *L*₂ norms. The quantity γ is the corresponding γ_0 or γ_{DG} . From the results in this table, we can conclude that in the $\|\cdot\|_{DG}$ norm the dominating error is the interpolation error, and as the next example shows, in general, the order of convergence in $\|\cdot\|_{DG}$ is 1.

The second test is for a fixed right hand side $\mathbf{f} = 2(1, x)$. We calculate approximations to the order of convergence of the numerical solutions on successively refined grids as follows:

$$\begin{aligned} \gamma &= \log_2 \frac{\|\mathbf{u}_k - \mathbf{u}_{k-1}\|}{\|\mathbf{u}_{k+1} - \mathbf{u}_k\|}, & \gamma_* &= \log_2 \frac{|\llbracket \mathbf{u}_k \rrbracket|_* - |\llbracket \mathbf{u}_{k-1} \rrbracket|_*}{|\llbracket \mathbf{u}_{k+1} \rrbracket|_* - |\llbracket \mathbf{u}_k \rrbracket|_*}, \\ \gamma_p &= \log_2 \frac{\|p_k - p_{k-1}\|_{0,\Omega}}{\|p_{k+1} - p_k\|_{0,\Omega}}, & k &= 1, \dots, 4. \end{aligned}$$

Square domain					L -shaped domain				
k	1	2	3	4	k	1	2	3	4
γ_0	1.77	1.86	1.93	1.97	γ_0	1.28	1.44	1.56	1.50
γ_{DG}	0.87	0.92	0.97	0.98	γ_{DG}	0.53	0.46	0.44	0.40
γ_p	0.98	0.99	0.99	1.0	γ_p	0.97	0.93	0.87	0.75
γ_*	0.90	0.95	0.98	0.99	γ_*	0.40	0.40	0.37	0.35

Table 6.2. Approximate order of convergence of the error for square and L -shaped domains and right-hand side $\mathbf{f} = 2(1, x)$.

Again, $\|\cdot\|$ denotes any of the (semi)-norms of interest and γ approximates the corresponding order of convergence. Table 6.2 shows the tabulated values of γ_0 , γ_{DG} , γ_p , and γ_* . It is clear from these values that the order of approximation for the velocity and the pressure is optimal for the square domain, whereas for the L -shaped domain the convergence is not of optimal order, due to the singularity of the solution near the reentrant corner.

6.3. Uniform preconditioning. The tests presented in this subsection illustrate the efficient solution of the system (6.3) below by Preconditioned Conjugate Gradient (PCG) with the preconditioner given in (6.4). We introduce the matrices representing the bilinear forms defined in (3.6)–(3.7), and also the mass matrix for the $\mathbf{BDM}_1$ space. We denote by $\mathbf{M}$ the mass matrix on $\mathbf{V}_h$ and by $\tilde{\mathbf{A}}$ the stiffness matrix associated with $a_h(\cdot, \cdot)$ on $\mathbf{V}_h$ in (3.6)–(3.7). We note that $\mathbf{A}$, without the divergence-free constraint, is spectrally equivalent to two scalar Laplacians.

It is known that the null space of $b(\cdot, \cdot)$ in (3.6) is made of vector fields that are curls of continuous, piecewise quadratic functions vanishing on the boundary. We denote by $\mathbf{P}_{\text{curl}}$ the matrix representation of the natural inclusion of these curls in the BDM space. Namely,

$$\mathbf{curl}(\text{basis functions in } \mathcal{N}_h) = (\text{basis functions in } \mathbf{V}_h) \mathbf{P}_{\text{curl}}.$$

It is easy to see that

$$\mathbf{A}_q = \mathbf{P}_{\text{curl}}^T \mathbf{M} \mathbf{P}_{\text{curl}}.$$

where $\mathbf{A}_q$ is the discretization of the Laplacian on N_h with homogeneous Dirichlet boundary conditions.

The problem of finding the solution of (5.5) then amounts to solving the following algebraic system of equations

$$(6.3) \quad \mathbf{P}_{\text{curl}}^T \tilde{\mathbf{A}} \mathbf{P}_{\text{curl}} \mathbf{U} = \mathbf{P}_{\text{curl}}^T \mathbf{F}.$$

Here the superscript T means that the adjoint is taken with respect to the ℓ_2 -inner product, $\mathbf{U}$ is the vector containing the velocity degrees of freedom, and $\mathbf{F}$ is the vector representing the right-hand side $(\mathbf{f}, \mathbf{v})$ of the problem (3.6).

The matrix representation $\mathbf{B}$ of the preconditioner B described in the previous section has the following form:

$$(6.4) \quad \mathbf{B} = \mathbf{A}_q^{-1} \mathbf{P}_{\text{curl}}^T \mathbf{M} \tilde{\mathbf{B}} \mathbf{M} \mathbf{P}_{\text{curl}} \mathbf{A}_q^{-1}$$

where $\tilde{\mathbf{B}}$ is a preconditioner of $\tilde{\mathbf{A}}$. In summary, the action of the preconditioner requires the solution of systems corresponding to 4 scalar Laplacians. It is also worth noting that suitable multigrid packages for performing these tasks are available today.

The results are summarized in Table 6.3. The legend for the symbols used in the table is as follows: n_{it} is the number of PCG iterations; ρ is the average reduction per one such iteration defined as $\rho = \left[\frac{\|r_{n_{it}}\|_{\ell_2}}{\|r_0\|_{\ell_2}} \right]^{1/n_{it}}$; J is the refinement level, for which $h \approx 2^{-J} h_0$, where h_0 is the characteristic mesh size on the coarsest grid. From the results in Table 6.3, we

Square domain						
J	0	1	2	3	4	5
n_{it}	5	6	6	7	7	7
ρ	0.06	0.07	0.09	0.10	0.11	0.13

L -shaped domain						
J	0	1	2	3	4	5
n_{it}	6	7	8	9	9	10
ρ	0.08	0.10	0.13	0.18	0.21	0.21

Table 6.3. Preconditioning results for square domain (left) and L -shaped domain (right). The PCG iterations are terminated when the relative residual is smaller than 10^{-6} .

can conclude that the preconditioner is uniform with respect to the mesh size. It is also evident that this method is in fact quite efficient in terms of the number of iterations and the reduction factor.

To conclude this section, let us point out that the generalizations to the 3D version of the preconditioner, and the introduction of the approximate solvers in (6.4) can be achieved in a similar fashion and are the subject of ongoing research.

APPENDIX A. PROOF OF PROPOSITION 3.3

We now state and prove a result, Proposition A.1 given below, which contains as a particular case the Proposition 3.3, used in Section 3 to show Korn inequality (cf. Lemma 3.2) for the case $d = 2$. After giving its proof, we comment briefly on how the result can be applied to show the corresponding Korn inequality (3.13) (cf. Lemma 3.2) for $d = 3$.

Proposition A.1. *Let T be a triangle (or a tetrahedron for $d = 3$) with minimum angle $\theta > 0$, and let e be an edge (resp. face) of T . Then for every $p > 2$ and for every integer k_{max} a constant $C_{p,\theta,k_{max}}$ exists such that*

$$(A.1) \quad \int_e \mathbf{v} \cdot (\boldsymbol{\tau} \cdot \mathbf{n}) \, ds \leq C_{p,\theta,k_{max}} h_T^{-1/2} \|\mathbf{v}\|_{0,e} h_T^{\frac{d(p-2)}{2p}} \|\boldsymbol{\tau}\|_{0,p,T}$$

for every $\boldsymbol{\tau} \in (L^p(\Omega))_{sym}^{d \times d}$ having zero divergence and for every $\mathbf{v} \in \mathbf{P}^{k_{max}}(T)$.

Proof. First we go to the reference element $\hat{T}$:

$$(A.2) \quad \left| \int_e \mathbf{v} \cdot (\boldsymbol{\tau} \cdot \mathbf{n}) \, ds \leq C_\theta |e| \left| \int_{\hat{e}} \hat{\mathbf{v}} \cdot (\hat{\boldsymbol{\tau}} \cdot \hat{\mathbf{n}}) \, d\hat{s} \right| \leq C_\theta h_e^{d-1} \left| \int_{\hat{e}} \hat{\mathbf{v}} \cdot (\hat{\boldsymbol{\tau}} \cdot \hat{\mathbf{n}}) \, d\hat{s} \right|$$

where $\hat{\mathbf{v}}$ and $\hat{\boldsymbol{\tau}}$ are the usual covariant and contravariant images of $\mathbf{v}$ and $\boldsymbol{\tau}$, respectively. And, here and throughout his proof, the constants C_θ and $C_{\theta, k_{max}}$ may assume different values at different occurrences. Note that the divergence-free property of $\boldsymbol{\tau}$ will be preserved by the contravariant mapping, and that $\mathbf{v}$ will still be a vector-valued polynomial of degree $\leq k_{max}$. Then for every component $\hat{v}$ of $\hat{\mathbf{v}}$, we construct the auxiliary function φ_v as follows. First we define φ_v on $\partial\hat{T}$ by setting it as equal to $\hat{v}$ on $\hat{e}$ and zero on the rest of $\partial\hat{T}$. Then we define φ_v in the interior using the harmonic extension. It is clear that φ_v will belong to $W^{1,p'}(\hat{T})$ (remember that $p > 2$ so that its conjugate index p' will be smaller than 2). Using the fact that $\hat{\mathbf{v}}$ is a polynomial of degree $\leq k_{max}$, it is not difficult to see that

$$(A.3) \quad \|\varphi_v\|_{W^{1,p'}(\hat{T})} \leq \hat{C}_{\theta, k_{max}} \|\hat{\mathbf{v}}\|_{0, \hat{e}}.$$

A simple integration by parts (remembering that $\text{div } \boldsymbol{\tau} = 0$) now gives

$$(A.4) \quad \int_{\hat{e}} \hat{\mathbf{v}} \cdot (\hat{\boldsymbol{\tau}} \cdot \hat{\mathbf{n}}) \, d\hat{s} = \int_{\partial\hat{T}} \varphi_v \cdot (\hat{\boldsymbol{\tau}} \cdot \hat{\mathbf{n}}) \, d\hat{s} = \int_{\hat{T}} \nabla \varphi_v : \hat{\boldsymbol{\tau}} \, d\hat{x} \\ \leq \|\varphi_v\|_{W^{1,p'}(\hat{T})} \|\hat{\boldsymbol{\tau}}\|_{(L^p(\hat{T}))_{sym}^{d \times d}} \leq \hat{C}_{\theta, k_{max}} \|\hat{\mathbf{v}}\|_{0, \hat{e}} \|\hat{\boldsymbol{\tau}}\|_{(L^p(\hat{T}))_{sym}^{d \times d}}.$$

The proof then follows immediately by inserting (A.4) into (A.2) and then coming back to the current element T by means of:

$$(A.5) \quad \|\hat{\mathbf{v}}\|_{0, \hat{e}} \leq C_\theta h_e^{-\frac{d-1}{2}} \|\mathbf{v}\|_{0, e} \quad \|\hat{\boldsymbol{\tau}}\|_{(L^p(\hat{T}))_{sym}^{d \times d}} \leq C_\theta h_T^{-\frac{d}{p}} \|\boldsymbol{\tau}\|_{(L^p(T))_{sym}^{d \times d}}$$

□

With this result in hand, we can show the Korn inequality (3.13) given in Lemma 3.2 for $d = 3$. It is necessary to modify the proof in only two places: the definition of the space of rigid motions on Ω , $\mathbf{RM}(\Omega)$, and the application of Proposition 3.19. The space $\mathbf{RM}(\Omega)$ is now defined by:

$$\mathbf{RM}(\Omega) = \{ \mathbf{a} + \mathbf{b}\mathbf{x} : \mathbf{a} \in \mathbb{R}^d \quad \mathbf{b} \in so(d) \}$$

with $so(d)$ denoting the Lie Algebra of the skew-symmetric $d \times d$ matrices.

To prove (3.16) (and so conclude the proof of (3.13)), estimate (3.21) is replaced by estimate (A.6) below, which is obtained as follows: first, by applying (A.1) (instead of (3.19)) from Proposition A.1 to each e in the last term in (3.20) and then by using the generalized Hölder inequality with the same exponents as for $d = 2$ (with $q = 1/2$ and

$r = 2p/(p-2)$, so that $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$)

$$(A.6) \quad \sum_{e \in \mathcal{E}_h^o} \int_e \llbracket \mathbf{v}_t \rrbracket : \{\boldsymbol{\tau}\} \, ds \leq C \left(\sum_{e \in \mathcal{E}_h^o} h_e^{-1} \|\mathbf{v}_t\|_{0,e}^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}_h^o} \|\boldsymbol{\tau}\|_{0,p,T(e)}^p \right)^{1/p} \left(\sum_{e \in \mathcal{E}_h^o} h_e^{\frac{d(p-2)}{2p}r} \right)^{1/r} \\ \leq C \|\mathbf{v}_t\|_*^2 \|\boldsymbol{\tau}\|_{0,p,\Omega} \mu(\Omega)^{1/r}.$$

Here, as in estimate (3.21), $T(e)$ in the first line denotes the set of elements T whose boundary contains e , $\mu(\Omega)$ in the second line denotes the measure of the domain Ω , and the constant C still depends on p , k_{max} , and on the maximum angle in the decomposition $\mathcal{T}_h$. The rest of the proof of Lemma 3.2 proceeds as for $d = 2$.

ACKNOWLEDGMENTS

Part of this work was completed while the first author was visiting IMATI-CNR of Pavia. She is grateful to the IMATI for the kind hospitality. The first author was partially supported by MEC through the project MTM2011-27739-C04-04. The second and third authors were partially supported by the Italian MIUR through the project PRIN2008. The last two authors were partially supported by US Department of Energy through the project de-sc0009603.

REFERENCES

- [1] R. A. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] S. Agmon. *Lectures on elliptic boundary value problems*. Prepared for publication by B. Frank Jones, Jr. with the assistance of George W. Batten, Jr. Van Nostrand Mathematical Studies, No. 2. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto-London, 1965.
- [3] D. N. Arnold, F. Brezzi, B. Cockburn, and L. D. Marini. Unified analysis of discontinuous Galerkin methods for elliptic problems. *SIAM J. Numer. Anal.*, 39(5):1749–1779 (electronic), 2001/02.
- [4] D. N. Arnold, F. Brezzi, and L. D. Marini. A family of discontinuous Galerkin finite elements for the Reissner-Mindlin plate. *J. Sci. Comput.*, 22/23:25–45, 2005.
- [5] D. N. Arnold, R. S. Falk, and R. Winther. Finite element exterior calculus, homological techniques, and applications. *Acta Numer.*, 15:1–155, 2006.
- [6] S. C. Brenner. Poincaré-Friedrichs inequalities for piecewise H^1 functions. *SIAM J. Numer. Anal.*, 41(1):306–324 (electronic), 2003.
- [7] F. Brezzi, J. Douglas, Jr., M. Fortin, and L. D. Marini. Efficient rectangular mixed finite elements in two and three space variables. *RAIRO Modél. Math. Anal. Numér.*, 21(4):581–604, 1987.
- [8] F. Brezzi, J. Douglas, Jr., and L. D. Marini. Two families of mixed finite elements for second order elliptic problems. *Numer. Math.*, 47(2):217–235, 1985.
- [9] F. Brezzi and M. Fortin. *Mixed and hybrid finite element methods*, volume 15 of *Springer Series in Computational Mathematics*. Springer-Verlag, New York, 1991.
- [10] B. Cockburn, G. Kanschat, and D. Schötzau. A note on discontinuous Galerkin divergence-free solutions of the Navier-Stokes equations. *J. Sci. Comput.*, 31(1-2):61–73, 2007.

- [11] H. C. Elman, D. J. Silvester, and A. J. Wathen. *Finite elements and fast iterative solvers: with applications in incompressible fluid dynamics*. Numerical Mathematics and Scientific Computation. Oxford University Press, New York, 2005.
- [12] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations. Vol. I*, volume 38 of *Springer Tracts in Natural Philosophy*. Springer-Verlag, New York, 1994. Linearized steady problems.
- [13] V. Girault and P.-A. Raviart. *Finite element methods for Navier-Stokes equations*, volume 5 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 1986. Theory and algorithms.
- [14] C. Greif, D. Li, D. Schötzau, and X. Wei. A mixed finite element method with exactly divergence-free velocities for incompressible magnetohydrodynamics. *Comput. Methods Appl. Mech. Engrg.*, 199(45-48):2840–2855, 2010.
- [15] O. A. Ladyzhenskaya. *The mathematical theory of viscous incompressible flow*. Second English edition, revised and enlarged. Translated from the Russian by Richard A. Silverman and John Chu. Mathematics and its Applications, Vol. 2. Gordon and Breach Science Publishers, New York, 1969.
- [16] S. V. Nepomnyaschikh. Mesh theorems on traces, normalizations of function traces and their inversion. *Soviet J. Numer. Anal. Math. Modelling*, 6(3):223–242, 1991.
- [17] S. V. Nepomnyaschikh. Decomposition and fictitious domains methods for elliptic boundary value problems. In *Fifth International Symposium on Domain Decomposition Methods for Partial Differential Equations (Norfolk, VA, 1991)*, pages 62–72. SIAM, Philadelphia, PA, 1992.
- [18] P. Oswald. Preconditioners for nonconforming discretizations. *Math. Comp.*, 65(215):923–941, 1996.
- [19] P.-A. Raviart and J. M. Thomas. A mixed finite element method for 2nd order elliptic problems. In *Mathematical aspects of finite element methods (Proc. Conf., Consiglio Naz. delle Ricerche (C.N.R.), Rome, 1975)*, pages 292–315. Lecture Notes in Math., Vol. 606. Springer, Berlin, 1977.
- [20] J. Xu. The auxiliary space method and optimal multigrid preconditioning techniques for unstructured grids. *Computing*, 56(3):215–235, 1996. International GAMM-Workshop on Multi-level Methods (Meisendorf, 1994).

CENTRE DE RECERCA MATEMÀTICA, UAB SCIENCE FACULTY, 08193 BELLATERRA, BARCELONA, SPAIN

IUSS-PAVIA C/P IMATI-CNR, VIA FERRATA 5/A 27100 PAVIA ITALY, AND DEPT. OF MATH., KAU, PO BOX 80203, JEDDAH, 21589, SAUDI ARABIA

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK PA 16802, USA

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK PA 16802, USA